
SCAMIT CODE: LACO 68 (in part) Date Examined: Feb. 10, 1986
Voucher by: Douglas Diener

Literature: Lomakina, N.B. 1958.

Diagnostic Characters:

1. Carapace with a single shallow lateral sulcus (check carapace sculpturing carefully with figure).
2. Pigment lines around sulcus are raised ridges.
3. Inner margin uropodal peduncle lacks spines but has short setae.

Related Species and Character Differences:

Campylaspis sp. B - anteriodorsal surface of carapace smooth, with shallow pits, unlike Campylaspis nr. crisper which has small pustules (e.g. like C. rubromaculata); raised ridges around lateral sulcus are absent in Campylaspis sp. B.

Comments: C. nr. crisper differs from C. crisper Lomakina 1955 by having an antennal notch and pseudorostrum; uropods are more slender with rami approximately 2/3 the length of the peduncle; rostral notch shallow and defined at ventral edge by small spine; inner margin of uropod inner ramus with four long spines (communicated by Don Cadien, MBC).

Distribution: Pt. Estero to Coronado Canyon; 118-610 m.





Figure 1. Male carapace (from Cadien, unpublished)

Figure 2. Male telson (from Cadien, unpublished)

Figure 3. Female telson; inner margin of uropod peduncle with 8 spines, inner margin of endopodite with 4 spines (from Phillips, unpublished).