

Table 8. Holothuroid distribution in the Central Eastern Pacific. Continued.

Taxon	NE P	32°N	26°N	20°N	14°N	8°N	2°N	4°S	10°S	SE P	0°E	10	20	110	200	1100	2000	4000	6000	Substrate
5114 <u>Pannychia moseleyi</u>		*--**--*--*--*--*--*	***_												212	2599				Ooze/glb gy;Md/gr Sand
5115 <u>Psychronaetes hanseni</u>					???												4800-5200			Oz/di;Mg Clay
FAMILY PSYCHROPOTIDAE																				
5116-5121																				
5116 <u>Benthodytes incerta</u>					*-----*												2417-4087			Mud/stky gy;O/glb
5117 <u>B. sanguinolenta</u>		**--*--*--*--*--*--*														(768)1975		7250		Md;Cl;Oz glb;Rhab
5118 <u>B. typica</u>					*-----*										315			4700		Ooze/glb gy
5119 <u>Psychropotes depressa</u>					*-----*											957		4060		Ooze/glb gr;Mud
5120 <u>P. longicauda</u>					*-----*													2210-5203		Oz/di,gr Md/fn;Mg
5121 <u>P. verrucosa</u>					**													2417	7250	Ooze/glb gr,yl
		S, B & G Cal	Mex	C Am	Pan-G-Ec	Peru						shallow	shelf			slope		abyss		