

SYNONYMY: Enopla sp 2 Pasko 2019

LITERATURE:

- Coe, W.R. 1905. Nemerteans of the west and northwest coasts of America. Bulletin of the Museum of Comparative Zoology, 47: 1–319
- Coe, W.R. 1940. Revision of the Nemertea fauna of the Pacific coasts of North, Central and northern South America. Allan Hancock Pac. Exped. 2(13): 247–323.

DIAGNOSTIC CHARACTERS (from a single specimen, Bight'18, Station 10438, 3.3m)

- Fairly elongate, thin species, ~8mm long, with smooth, cream colored epidermis
- Eyes 3 pairs, anterior set of four form a square, while third pair is slight more distant from others
- Proboscis sheath approximately $\frac{1}{2}$ the length of the body
- Stylet apparatus located mid-body; stylet subequal to basis; basis notably thin, slightly tapered in mid-region with narrowly rounded base
- Two accessory pouches present, each with a single accessory stylet


Similarities:

- Differs from all currently recognized enoplans in the fairly distinctive eye pattern and stylet apparatus.

Hoplonemertea sp DC2,
dorsal view, clear




Hoplonemertea sp DC2,
eye pattern



Hoplonemertea sp DC2,
Stylet apparatus

1mm



Hoplonemertea sp
DC2, dorsal view