

Related Species:

Similar to Japanese S. mourense except for presence of 2-4 fusiform bodies, rathe rthan 5.

Comments:

Longitudinal muscle bands are most visible with light shining through the base of specimen.

References:

- Cutler, Edward Bayler. <u>The Sipuncula: their systematics, biology and evolution</u>. Ithaca: Comstock Publishing Associates, 1994.
- Stephen, A.C. & S.J. Edmonds. <u>The Phyla Sipuncula and Echiura</u>. London: The British Museum (Natural History), 1972.

Species: *Siphonosoma ingens*

Taxon: Sipunculida: Sipunculidae Date: November 15, 1995

Additional Illustrations: (Fisher, 1947)

Fig. 2

Head of a small specimen (Monterey Bay, CA)




Fig. 3

Portion of Fig. 1 enlarged

CP, coelomic papillae; CV, dorsal contractile vessel; N, nephridium; NC, nerve cord; O, esophagus; RD, dorsal retractor muscle; RV, ventral retractors.





Fig. 4

Outer body wall musculature of *S. ingens;* half body width with 4 muscle bands shown.



Distribution— (Cutler, 1994) Pt. Loma: 200 ft.

Geographic: California

Habitat: intertidal to subtidal, sand to sandy mud