

SCAMIT Code: LACO 82

Date examined: October 11, 1988
Voucher by: Tom Parker

Synonymy: Lumbrineris bifilaris Hartman, 1944, 1968
(not Ehlers, 1901)

Literature:

Fauchald, K. 1970. Polychaetous Annelids of the Families Eunicidae, Lumbrineridae, Iphitimidae, Arabellidae, Lysaretidae, and Dorvilleidae from Western Mexico. Allan Hancock Monograph in Marine Biology. No. 5. 335 p.

Hartman, O. 1944. Polychaetous Annelids. Pt. 5. Eunicea. Allan Hancock Pacific Expeditions. 10: 1-238.

Hartman, O. 1968. Atlas of the Errantiate Polychaetous Annelids From California. Allan Hancock Foundation. 828 p.

Diagnostic Characters: *fits some of "genotype"*

1. Acicula yellow.
2. Simple hooded hooks present from first setiger.
3. Posterior presetal and postsetal lobes prolonged.
4. Maxillary formula: 1-(4-5)-2-1.

Comments: The description in Hartman's Atlas (1968) incorrectly states that the acicula are black.

Distribution: Throughout southern California, in shelf depths.




Fig. 1 Posterior parapod and lobes



Fig. 2 Maxilla

(Figures from Hartman, 1968)