

only complete

8  
7

Lumbrineris latreilli Audouin and Milne Edwards, 1934  
Lumbrineridae

Vol. 7, No. 7

SCAMIT Code: HYP 79

Date examined: October 11, 1988

Voucher by: Tom Parker

needs a number

Synonymy: None

Literature:

Fauchald, K. 1970. Polychaetous Annelids of the Families Eunicidae, Lumbrineridae, Iphitimidae, Arabellidae, Lysaretidae, and Dorvilleidae from Western Mexico. Allan Hancock Monograph in Marine Biology. No. 5. 335 p.

Hartman, O. 1944. Polychaetous Annelids. Pt. 5. Eunicea. Allan Hancock Pacific Expeditions. 10: 1-238.

Hartman, O. 1968. Atlas of the Errantiate Polychaetous Annelids From California. Allan Hancock Foundation. 828 p.


Diagnostic Characters: *SUMMIT MOUNT*

*per Lewis*

1. Acicula yellow.
2. Composite hooded hooks present in anterior setigers; the last appearing on setiger 11-24.
3. Presetal and postsetal lobes not prolonged in any setiger.
4. Maxillary formula: 1-5-2-1.

Comments: Acicula color is clearly different from acicula in L. japonica.

Distribution: Throughout southern California, in shelf depths. From Santa Monica Bay in 18-45 m in mixed sediments.



Maxilla (from Hartman, 1968)