SCAMIT Code: OC 65

Date Examined: 14 April 1986 Voucher By: Larry Lovell(MEC)

Literature:

Banse, 1972 (syn. <u>C</u>. <u>ecaudata</u>) Hartman, 1944 Hartman, 1969 Rowe, 1980

Diagnostic Characters:

- A short tumid species of Chone.
- Collar stains lightly with a ventral unstained cresent shaped area just anterior of setiger one, no stain on the collar dorsally.
- Ventral shields present.

Related species and Differences:

- 1. <u>C</u>. <u>ecaudata</u> is a slimmer bodied animal and is found in finer sediments.
- 2. <u>C.</u> sp. B (<u>sensu Harris</u>) is very small (<u>Euchone</u> sized) and has very dark lateral staining on the collar and setiger one.
- 3. All other species of <u>Chone</u> from California lack ventral shields.

Additional Remarks:

Banse, 1972 considered \underline{C} . minuta a synonym of \underline{C} . ecaudata, but local taxonomists consider them as separate species. \underline{C} . minuta is found in rocky and mixed sediment, while \underline{C} . ecaudata is found in mud and sand.

Distribution:

Baja to Pt. Conception, intertidal to 60m.




Figure a. Anterior end, ventral view showing methyl green staining pattern.