

SCAMIT Code: Provisional

Date examined: June 12, 1986

Voucher by: Leslie H. Harris
(MBC)

SYNONYMY: Jasmineira sp.A Harris
Fabrisabella sp.A Williams

LITERATURE: Annenkova 1937
Banse 1979
Fauvel 1927

DIAGNOSTIC CHARACTERS:

1. Body long and linear, 7mm without crown.
8 thoracic and about 25 setigers.
2. Thoracic notosetae 3 kinds: bilimbate, subspatulate and spatulate with short mucron. Abdominal neurosetae capillary or unilimbate with very narrow wings.
3. Thoracic neurosetae long-handled, acicular uncini, gently bent, 7-12 per fascicle. Abdominal uncini avicular, S-shaped, strongly-bent, 10-11 per fascicle. Both types have large main fang and crest of smaller teeth. No companion setae.
4. Collar with ventral notch, fairly level all the way around, dips slightly dorsally. Dorsal edges infolded, rarely gaps, brancial basis covered.
5. Anus ventral, pygidium has digitate caudal appendage. No eyespots.
6. Otocysts (vascular folds) present, 1 pair, dorsally within the collar.
7. 10-12 pairs of radioles, long. Branchial crown usually broken off slightly above branchial basis. No flange or palmate membrane (?).
8. Narrow glandular girdle on setiger 2. Intrasegmental furrows present in thoracic setigers.