

# *Eohaustorius estuarius*

## Diagnostic Characters:

Rostrum moderate to weak  
Coxa 1-4 enlarging, coxa 4 dominate  
Gnathopods feeble and simple  
Pereopod 4 smaller and unlike pereopod 3, reversed in orientation

Peraeopod 7, segment 6 with 2 groups of posterior marginal spine groups

Coxae 3 & 4 antero-distal margins rounded

P5, segment 4 less broad, 1.5 X length

Pleon overhang normal, little or not exceeding side plate process

Uropod 3, inner ramus with 5 marginal setae

P4, segment 5, hind lobe normal, width about equal to length

P7, segment 6 with 3-4 posterior marginal spines groups

G2, basis, hind margin setose throughout

P4, segment 5, posterior lobe, distal margin with spine cluster

P6, segment 4 with lower facial row of 2-3 spine groups

P6, segment 5 with posterior marginal spine group

P6 segment 4 subtriangular, widest distally, length <2X width; basis as broad as length

## Illustration:

AMPHIPACIFICA VOL. 11 NO. 1, AUG. 31, 1995 42




FIG. 4. *Eohaustorius estuarius* Bosworth, Female ov (5.0 mm) Long Beach, V. I., B. C. (partly after Bosworth, 1973)