

Tabular Guide to Female Cylindroleberididae reported by the major monitoring programs  
in the Southern California Bight

Character	<i>Xenoleberis</i>	<i>Bathyleberis</i>	<i>Bathyleberis</i>	<i>Bathyleberis</i>	<i>Postasterope</i>	<i>Postasterope</i>	<i>New Genus</i>
	<i>californica</i>	<i>sp.</i>	<i>garthi</i>	<i>hancocki</i>	<i>barnesi</i>	<i>hulingsi</i>	<i>SD1</i>
Carapace Lenth	2.66 - 2.69	2.20 - 2.60	1.80 - 1.85	2.17 - 2.59	1.43 - 1.59	1.34 - 1.60	2
Triaenid & spinose bristles on endite of mandible	3/3	3/4	3/3	3/4	3/4	2/3	
Bristles proximal to a-bristle of mandible	2	2-3	2	3	1	2	1
Bristles between a- and b-bristle of mandible	1	2	1	2	0	0	1
Bristles lateral to b- bristle of mandible	0	3-4	3	6	0	0	1
Length of exopodite in relation to 1st endopodite podomere of mandible	1/10	1/2 - 2/3	1/2	2/3	2/3	1/2	1/2
Bristles along ventral margin of 6th limb	19	15 - 17	12	20	3	2	15 - 16
Lateral bristles of 7th limb	6	6 - 9	6	11	6	4	9
Caudal furca claw pattern	10	9	10	10	8	9	10
Presence of e-bristle	no	yes	yes	yes	no	no	
Bristles lateral to c-bristle	6	6	6	6	6	5	
Accessory filament on sensory bristle	yes	yes	yes	yes	no	no	