

Ostracods From Southern California
of the
Genus Rutiderma

Brad Myers *

Subclass Ostracoda
Order Myodocopida
Suborder Myodocopina
Family Rutidermatidae

Rutiderma hartmani Foulson 1965

Range: Catalina Island- Panama

Depth: 9-57m

Rutiderma judayi McKenzie 1965

Range: Santa Rosa Island- Scammon Lagoon, Baja Calif.

Depth: Shallow sub-tidal- 13m

Rutiderma lomae (Juday 1907)

Synonymy: Philomedas lomae Juday 1907

Range: Oxnard, Calif- San Bartholome Bay, Baja Calif.

Depth: 11-401m

Rutiderma rostrata Juday 1907

Range: Ft. Conception- El Salvador

Depth: 5.2-97.5m

Rutiderma rotunda Foulson 1965 (June 10, 1965)

Synonymy: Rutiderma californica McKenzie 1965 (July 15, 1965)

Range: Santa Cruz Island, Calif- Guyamas, Mex.

Depth: Intertidal- 33.8m

Note-- Distribution data based on literature and personal records.

SCCWRP Taxonomic Standardization Program, February 1979

* 2951 "E" So. Fairview / Santa Ana, Calif. 92704 (714)545-6666

California
Anterior of Shell

derman
Lateral View




♀


R. rotunda




R. rostrata




R. judayi





R. lomae



~~R. lactum~~
R. chessi




♂




? unknown ?

California Idernan
 Posterior of Shell Lateral View



♀


R. rotunda^{var}




R. rostrata^{var}




R. judayi






R. lomae



~~R. hartmanni~~^{hi}
 R. chessi



♂



? unknown ?

