

Marnovich, 1977

*Calinaticina oldroydii*

(Dall, 1897)

Shell relatively large, thin; whorls 4; spire elevated; umbilicus narrow; parietal wall thin, edge straight, reflected to partially obstruct umbilicus; spiral sculpture of fine striae; axial sculpture of growth increments; light brown periostracum; periostracum shaggy within umbilicus; operculum corneous, paucispiral filling aperture. Length 40 - 70 mm. Soft bottoms, 30 - 400 m. (McLean, 1996)




Marnovich, 1977

*Neverita reclusiana*

(Deshayes, 1839)

Shell tan colored, whorls flat sided; a heavy white callus extending from the suture covers the umbilicus except for a chink. Height, 40 - 70 mm. (McLean, 1978)




Marnovich, 1977

*Polinices lewisii*

(Gould, 1847)

Shell large, heavy, with a shallow channel at the shoulder of the whorl, columellar callus extending to the side of the deep, narrow umbilicus. Height 75 - 130 mm. (McLean, 1978)



Marnovich, 1977

*Polinices draconis*

(Dall, 1903)

With a thinner shell than *P. lewisii*, and not quite as large, this species has rounded or tabulate shoulder, no columellar callus, and a broad umbilicus showing a thick periostracum. Height 50 - 90 mm. (McLean, 1978)



Marnovich, 1977

*Sinum scopulosum*

(Conrad, 1849)

Shell with a greatly inflated body whorl, depressed spire, and sculpture of fine incised spiral lines. Color, yellowish white under a fine periostracum.

Maximum diameter, 30 - 40 mm. (McLean, 1978)

