
City of San Diego **PROVISIONAL SPECIES VOUCHER SHEET**

Provisional Name: Amphiporus punctatulus	Taxon: Nemertea: Amphiporidae			Taxonomist: D. Pasko		
Authority: Coe 1905	Date: 12 Aug	ust 1999				
Common Synonyms:	Specimen(s):	STATION	DATE	<u>DEPTH</u>	LOCATION	VIAL#
	Bight98 - SD Bay	2225	27Jul98	3.6 m	CSD	



Related Species & Other Comments:

A. punctatulus stands out from other amphiporids by the short, bulbous basis and fairly distinctive eye pattern. It is distinguished from A. bimaculatus by the absence of pigment patches on the head, and the shape of the stylet, and the arrangement of the ocelli. In A. bimaculatus the basis is pear shaped and more elongate rather than short and bulbous as in A. punctatulus. A. bimaculatus has 10^{\pm} ocelli arranged in a line on each side of the head anterior to the cerebral sense organ, followed with a more medial, paired cluster (frequently located beneath the triangular pigment patches). In A. *punctatulus*, the paired lines of ocelli (only four in this specimen) are followed by two separate clusters of ocelli.

Coe (1905) reports the distribution from San Pedro - San Diego, however, in Coe (1940) the species is reported as being "known only from Isthmus Cove, Catalina Island, California" beneath stones within muddy sediments.

References:

Coe, W.R. 1905. Synopsis of North American Invertebrates. XXI. The Nemerteans. Part 1. Species occurring on the west and northwest coasts of North America. Amer. Natur. 39 (463):425-447.

Coe, W.R. 1940. Revision of the nemertean fauna of the Pacific coasts of North America, Central, and northern South America. Alan Han. Pac. Exp. Vol 2 (13). 323 pp. filename:I:\dean\A punctatulusvoucher.p65